

## Citizens [among those who required one of these tests: MRIs, CAT scans and angiographies, in a non-emergency situation in the last 12 months] who waited less than a month for a diagnostic test

Unit : %

	Year 2020
<b>Province</b>	
New Brunswick	52.3
<b>Health zones</b>	
Zone 1 - Moncton and South-East Area	52.1
Zone 2 - Fundy Shore and Saint John Area	54.1
Zone 3 - Fredericton and River Valley Area	53.6
Zone 4 - Madawaska and North-West Area	45.9
Zone 5 - Restigouche Area	59.0
Zone 6 - Bathurst and Acadian Peninsula Area	46.6
Zone 7 - Miramichi Area	45.9
<b>Communities</b>	
Bathurst, Beresford, Petit-Rocher Area	43.1
Bouctouche, Richibucto, Saint-Antoine Area	49.8
Campbellton, Atholville, Tide Head Area	58.0
Caraquet, Paquetville, Bertrand Area	53.7
Dalhousie, Balmoral, Belledune Area	47.5
Dieppe and Memramcook	44.1
Douglas, Saint Marys, Doaktown Area	42.1
Edmundston, Rivière-Verte, Lac Baker Area	39.1
Florenceville-Bristol, Woodstock, Wakefield Area	62.0
Fredericton	54.5

	<b>Year 2020</b>
Grand Bay-Westfield, Westfield, Greenwich Area	<b>44.2</b>
Grand Falls, Saint-Léonard, Drummond Area	<b>53.3</b>
Hillsborough, Riverside-Albert, Alma Area	<b>48.2</b>
Kedgwick, Saint-Quentin and Grimmer	<b>60.4</b>
Minto, Chipman, Cambridge-Narrows Area	<b>50.0</b>
Miramichi, Rogersville, Blackville Area	<b>44.7</b>
Moncton	<b>58.5</b>
Nackawic, McAdam, Canterbury Area	<b>38.3</b>
Neguac, Alnwick, Esgenoopetitj Area	<b>44.1</b>
New Maryland, Kingsclear, Lincoln Area	<b>51.4</b>
Oromocto, Gagetown, Fredericton Junction Area	<b>49.7</b>
Perth-Andover, Plaster Rock, Tobique Area	<b>57.4</b>
Quispamsis, Rothesay, Hampton Area	<b>49.1</b>
Riverview and Coverdale	<b>46.0</b>
Sackville, Dorchester, Port Elgin Area	<b>46.1</b>
Saint John, Simonds and Musquash	<b>61.1</b>
Salisbury and Petitcodiac	<b>56.4</b>
Shediac, Beaubassin East and Cap-Pelé	<b>48.5</b>
Shippagan, Lamèque, Inkerman Area	<b>40.5</b>
St. George, Grand Manan, Blacks Harbour Area	<b>53.9</b>
St. Stephen, Saint Andrews, Campobello Island Area	<b>42.0</b>
Sussex, Norton, Sussex Corner Area	<b>54.1</b>
Tracadie and Saint-Isidore	<b>36.2</b>

## Age

	Year 2020
18-64	48.8
65 and over	54.6
<b>Sex at birth</b>	
Female	47.0
Male	54.9
<b>Sexual identity</b>	
Minority sexual identity	50.4
Heterosexual	50.1
<b>Language</b>	
Preferred language of service: English	53.0
Preferred language of service: French	44.8
<b>Indigenous identity</b>	
Indigenous	52.2
Non-Indigenous	50.3
<b>Immigrant</b>	
Immigrants	50.2
Non-immigrants	51.3
<b>Education level</b>	
High school or less	53.3
Postsecondary	49.1
<b>Household income</b>	
Less than \$25,000	57.2
\$25,000 to less than \$60,000	51.8

	Year 2020
\$60,000 or more	46.4
<b>Persons with a disability</b>	
Without disability	52.4
With disability	51.0

## About

This indicator is the percentage of citizens [among those who required one of these tests: MRIs, CAT scans and angiographies, in a non-emergency situation in the last 12 months] who waited less than a month for a diagnostic test.

## Source

Primary Health Survey and Primary Care Survey

## Calculations

This indicator is based on respondents who have required one of these tests: MRIs, CAT-scans or angiographies provided in a non-emergency situation in the last 12 months . This indicator is calculated by dividing the number of respondents who answered "less than one month" by the number of respondents who answered "less than one month, 1 to 3 months or longer than 3 months" to the question: How long did you have to wait between when you and your doctor or nurse decided to go ahead with the test and the day of the test? If you have had more than one in the last 12 months, please answer with the most recent in mind.

## Notes

The 2020 edition of the Primary Health Survey (PHS) was conducted during the COVID-19 pandemic. While the pandemic did not influence the survey's response rate, NBHC had to assess the impact of surveying during the pandemic. Accordingly, advanced statistical analyses were performed to allow for comparison between the results obtained in 2020 and those obtained in previous PHS cycles. Despite these analyses, users are advised to use the PHS 2020 data with caution, especially when creating estimates for small sub-populations or when comparing it to other PHS editions.

---

### Unit

%

### Interpretation

Higher is better

### NBHC code

PH\_ACCOE\_1

### Caption

n/a = Not applicable / not available

S = Data suppressed due to confidentiality requirements and/or small sample size