

## Citizens who reported a health problem or long-term condition that has lasted or is expected to last for six months or more [among those who reported being limited sometimes or often in the kinds or amount of activity they can do at home, work or otherwis

Unit : %

	Year 2020
<b>Province</b>	
New Brunswick	<b>78.5</b>
<b>Health zones</b>	
Zone 1 - Moncton and South-East Area	<b>77.1</b>
Zone 2 - Fundy Shore and Saint John Area	<b>85.4</b>
Zone 3 - Fredericton and River Valley Area	<b>82.6</b>
Zone 4 - Madawaska and North-West Area	<b>75.9</b>
Zone 5 - Restigouche Area	<b>81.1</b>
Zone 6 - Bathurst and Acadian Peninsula Area	<b>68.8</b>
Zone 7 - Miramichi Area	<b>77.6</b>
<b>Communities</b>	
Bathurst, Beresford, Petit-Rocher Area	<b>65.9</b>
Bouctouche, Richibucto, Saint-Antoine Area	<b>74.9</b>
Campbellton, Atholville, Tide Head Area	<b>64.3</b>
Caraquet, Paquetville, Bertrand Area	<b>61.9</b>
Dalhousie, Balmoral, Belledune Area	<b>83.8</b>
Dieppe and Memramcook	<b>70.5</b>
Douglas, Saint Marys, Doaktown Area	<b>66.2</b>
Edmundston, Rivière-Verte, Lac Baker Area	<b>76.2</b>
Florenceville-Bristol, Woodstock, Wakefield Area	<b>80.7</b>

	Year 2020
Fredericton	<b>70.5</b>
Grand Bay-Westfield, Westfield, Greenwich Area	<b>83.3</b>
Grand Falls, Saint-Léonard, Drummond Area	<b>75.7</b>
Hillsborough, Riverside-Albert, Alma Area	<b>74.1</b>
Kedgwick, Saint-Quentin and Grimmer	<b>75.2</b>
Minto, Chipman, Cambridge-Narrows Area	<b>71.3</b>
Miramichi, Rogersville, Blackville Area	<b>69.2</b>
Moncton	<b>81.9</b>
Nackawic, McAdam, Canterbury Area	<b>72.2</b>
Neguac, Alnwick, Esgenoopetitj Area	<b>54.6</b>
New Maryland, Kingsclear, Lincoln Area	<b>82.9</b>
Oromocto, Gagetown, Fredericton Junction Area	<b>80.8</b>
Perth-Andover, Plaster Rock, Tobique Area	<b>78.3</b>
Quispamsis, Rothesay, Hampton Area	<b>82.8</b>
Riverview and Coverdale	<b>80.4</b>
Sackville, Dorchester, Port Elgin Area	<b>86.4</b>
Saint John, Simonds and Musquash	<b>83.2</b>
Salisbury and Petitcodiac	<b>90.2</b>
Shediac, Beaubassin East and Cap-Pelé	<b>62.6</b>
Shippagan, Lamèque, Inkerman Area	<b>54.3</b>
St. George, Grand Manan, Blacks Harbour Area	<b>73.2</b>
St. Stephen, Saint Andrews, Campobello Island Area	<b>77.5</b>
Sussex, Norton, Sussex Corner Area	<b>96.7</b>
Tracadie and Saint-Isidore	<b>69.6</b>

	Year 2020
<b>Age</b>	
18-64	75.0
65 and over	76.1
<b>Sex at birth</b>	
Female	76.7
Male	73.9
<b>Sexual identity</b>	
Minority sexual identity	76.0
Heterosexual	73.4
<b>Language</b>	
Preferred language of service: English	76.8
Preferred language of service: French	70.4
<b>Indigenous identity</b>	
Indigenous	76.2
Non-Indigenous	73.1
<b>Immigrant</b>	
Immigrants	67.6
Non-immigrants	78.0
<b>Education level</b>	
High school or less	73.9
Postsecondary	76.9
<b>Household income</b>	
Less than \$25,000	77.1

	Year 2020
\$25,000 to less than \$60,000	74.6
\$60,000 or more	78.5
<b>Persons with a disability</b>	
Without disability	n/a
With disability	78.5

## About

This indicator is the percentage of citizens who reported a health problem or long-term condition that has lasted or is expected to last for six months or more [among those who reported being limited sometimes or often in the kinds or amount of activity they can do at home, work or otherwise because of a physical or mental condition, or a health problem].

## Source

Primary Health Survey and Primary Care Survey

## Calculations

This indicator is based on respondents who have reported being limited . This indicator is calculated by dividing the number of respondents who answered "yes" by the number of respondents who answered "yes or no" to the question: What type of limitation or disability do you often or sometimes experience: k) Health problem or long-term condition that has lasted or is expected to last for six months or more

## Notes

The 2020 edition of the Primary Health Survey (PHS) was conducted during the COVID-19 pandemic. While the pandemic did not influence the survey's response rate, NBHC had to assess the impact of surveying during the pandemic. Accordingly, advanced statistical analyses were performed to allow for comparison between the results obtained in 2020 and those obtained in previous PHS cycles. Despite these analyses, users are advised to use the PHS 2020 data with caution, especially when creating estimates for small sub-populations or when comparing it to other PHS editions.

---

<b>Unit</b>	<b>Interpretation</b>	<b>NBHC code</b>
%	Lower is better	PH_PWDHP_1

## Caption

n/a = Not applicable / not available

S = Data suppressed due to confidentiality requirements and/or small sample size