

Tracadie-Sheila, Saumarez and Saint-Isidore

Canton des Basques, Haut-Sheila, Leech, Losier Settlement, Petit Tracadie, Pointe à Tom, Pointe des Robichaud, Pont Lafrance, Pont Landry, Rivière à la Truite, Saint-Irénée, Saint-Isidore, Saint-Pons, Saumarez, Tracadie Beach, Tracadie Beach, Saumarez, Tracadie-Sheila, Val-Comeau



Demographic Profile ¹		2014 (%)	NB (%)
Age Group	18-34	21.2	24.3
	35-54	39.3	36.7
	55-64	20.1	18.7
	65+	19.4	20.3
Gender	Male	49.3	48.0
	Female	50.7	52.0
Education Level	High school or less	44.0	38.7
	College/Technical diploma	32.0	31.4
	University degree	23.9	29.9
Aboriginal	Yes	7.1	3.0
Preferred Language of Service	English	4.0	74.6
	French	96.0	25.4
Household Income	Less than \$25,000	32.8	19.9
	\$25,000 - \$59,999	32.1	35.5
	\$60,000 or more	35.1	44.6
Employment Status	Employed	48.8	54.6
	Unemployed/Seasonal	10.1	7.3
	Retired	26.3	24.7
	Other	14.8	13.5

n = 240

n = 13,614

Population (18 and over) based on 2011 Census: 11,750

Confidence interval calculated at a 95% level of confidence.

Sample size shown represents total number of respondents.
Exact number of respondents varies per survey indicator.

Health Services Barriers	2011 (%)	2014 (%)	NB (%)
Health services not available in your area when needed	35.7 (30.7 – 40.8)	31.9 (26.0 – 37.8)	17.4 (16.7 – 18.0)
Transportation problems	6.0 ^E (3.5 – 8.4)	4.2 ^E (1.7 – 6.7)	7.1 (6.6 – 7.5)
Language problems with health care provider	13.7 (10.0 - 17.3)	4.1 ^E (1.6 – 6.5)	4.2 (3.9 – 4.6)
Cost for medication too high	68.8 (63.8 – 73.7)	61.1 (54.9 – 67.2)	35.2 (34.3 – 36.0)
Cost for treatments or procedures too high	42.7 (37.3 – 48.2)	41.9 (35.5 – 48.3)	19.9 (19.2 – 20.6)
Cost for ambulance services too high	59.6 (53.5 – 65.6)	37.0 (30.2 – 43.9)	21.7 (20.9 – 22.4)
Unable to leave the house because of health problems	19.6 (15.4 – 23.8)	11.0 ^E (7.0 – 14.9)	12.5 (11.9 – 13.0)
Have trouble finding their way around the health care system	12.1 (8.6 – 15.6)	7.0 ^E (3.8 - 10.3)	9.3 (8.8 – 9.8)
Difficulty understanding written information about medical condition or prescription (% always or usually)	17.6 (13.5 – 21.7)	10.4 ^E (6.5 – 14.2)	9.4 (8.9 – 9.9)
How often was a medical condition or prescription explained to you in a way that you could understand? (% always or usually)	86.1 (82.4 – 89.8)	87.2 (83.0 – 91.4)	91.0 (90.6 – 91.5)
Things that contribute a lot to feelings of stress			
Health of family members	Not Available	47.3 (41.0 – 53.5)	38.1 (37.3 - 38.9)
Time pressures / not enough time	Not Available	41.3 (35.1 – 47.5)	40.9 (40.1 – 41.7)
Your financial situation (ex: not enough money, or debt)	Not Available	34.7 (28.7 – 40.7)	33.5 (32.7 – 34.2)

n = 337

n = 240

n = 13,614

¹Based on survey respondents, weighted by age and gender

^E Use with caution (coefficient of variation between 16.6% and 33.3%)

^F Too unreliable to be published (coefficient of variation greater than 33.3%)

Tracadie-Sheila, Saumarez and Saint-Isidore

↑ = Higher than 2011 results
 ↓ = Lower than 2011 results

	n = 337	n = 240	n = 13,614
Health Profile	2011 (%)	2014 (%)	NB (%)
Self-rated Health (% very good or excellent)	39.7 (34.5–44.9)	32.0 (26.2–37.9)	50.5 (49.6–51.3)
Self-rated mental or emotional health (% very good or excellent)	Not Available	60.7 (54.6–66.9)	65.4 (64.6–66.2)
“My health largely depends on how well I take care of myself” (% strongly agree)	46.3 (41.0–51.6)	39.1 (33.0–45.2)	55.2 (54.4–56.0)
Number of prescription medications taken on a regular basis ³ (% 6 or more)	17.5 (12.7–22.3)	18.9 (13.1–24.8)	15.2 (14.5–15.9)
Discuss regularly with a health professional on improving health or preventing illness (% always or usually)	31.4 (26.2–36.6)	25.2 (19.8–30.7)	25.4 (24.7–26.1)
How long have you been seeing the same family doctor? (% 5 years or more)	74.2 (69.4–79.0)	69.9 (64.0–75.7)	73.0 (72.2–73.7)
Are you limited in doing activities because of a physical or mental condition, or a health problem? (% yes)	Not Available	23.6 (18.3–29.0)	22.1 (21.4–22.8)
Memory Loss			
Someone in household has a memory problem	Not Available	7.4 ^E (4.1–10.6)	9.9 (9.4–10.4)
...that interferes with day to day function	Not Available	F	3.9 (3.6–4.2)
Health Behaviours			
Eating fruits and vegetables (% 5 portions or more a day)	Not Available	48.7 (42.5–55.0)	50.4 (49.5–51.2)
Moderate or vigorous physical activity (% at least 2 ½ hours a week)	Not Available	46.4 (40.2–52.7)	49.0 (48.1–49.8)
Current smoker (% daily or occasional)	Not Available	19.0 (14.1–23.9)	19.2 (18.5–19.8)
Unhealthy weight (% obese)	Not Available	34.5 (28.5–40.6)	30.8 (30.0–31.6)

	n = 337	n = 240	n = 240	n = 13,614
Chronic Health Conditions ¹	2011 (%)	2014 (%)	2014 ² (#)	NB (%)
One or more chronic health conditions ³	65.6 (60.4–70.8)	70.9 (65.2–76.6)	8,329	61.6 (60.8–62.4)
High blood pressure	31.0 (26.1–35.9)	34.4 (28.5–40.3)	4,042	27.0 (33.3–38.3)
Arthritis	12.6 (9.1–16.1)	19.0 (14.1–23.9)	2,233	17.4 (16.8–18.0)
Gastric Reflux (GERD)	14.5 (10.8–18.2)	16.8 (12.2–21.5)	1,979	16.4 (15.8–17.0)
Chronic pain	15.6 (11.8–19.5)	16.4 (11.7–21.0)	1,922	14.0 (13.5–14.6)
Diabetes	13.4 (9.8–16.9)	16.1 (11.5–20.7)	1,892	10.7 (10.1–11.2)
Heart disease	9.8 (6.7–13.0)	15.1 (10.7–19.6)	1,780	8.3 (7.9–8.8)
Depression	13.4 (9.8–17.0)	15.0 (10.5–19.4)	1,758	14.9 (14.3–15.5)
Asthma	17.5 (13.5–21.4)	11.6 ^E (7.6–15.6)	1,361	11.8 (11.3–12.4)
Cancer	7.8 ^E (5.0–10.7)	5.7 ^E (2.8–8.6)	670	8.3 (7.8–8.7)
Mood disorder other than depression	F	F	357	3.0 (2.7–3.2)
Emphysema or COPD	3.8 ^E (1.7–5.8)	F	224	3.0 (2.7–3.3)
Stroke	3.8 ^E (1.8–5.8)	F	170	2.5 (2.2–2.8)

Confidence interval calculated at a 95% level of confidence.

Symbols ↑ and ↓ indicate a statistically significant difference at the 95% level of confidence.

Sample size shown represents total number of respondents. Exact number of respondents varies per survey indicator.

¹ Self-reported based on survey respondents

² Estimated number of citizens who have the selected chronic health condition

³ Among the twelve chronic health conditions listed in table located at the right of this page.

^E Use with caution (coefficient of variation between 16.6% and 33.3%)

^F Too unreliable to be published (coefficient of variation greater than 33.3%)

Tracadie-Sheila, Saumarez and Saint-Isidore

B = Better than 2011 results
W = Worse than 2011 results

n=337 n=240 1 to 33 n=13,614

Primary Health Indicator		2011	2014	Rank ²	NB		
Accessibility							
The ability of patients/clients to obtain care/service at the right place and the right time, based on respective needs, in the official language of their choice							
Personal Family Doctor	A-1	Has a personal family doctor (% yes)	91.5 (88.5 - 94.5)	96.6 (94.3 - 98.9)	6	92.1 (91.7 - 92.5)	
	A-2	Calling family doctor's office during regular practice hours (% very easy or somewhat easy)	71.9 (66.9 - 76.9)	77.0 (71.6 - 82.4)	21	78.3 (77.6 - 79.0)	
	A-3	Family doctor has extended office hours (after 5pm or on weekends) (% yes)	Not available	9.1^E (5.4 - 12.8)	28	16.2 (15.6 - 16.8)	
	A-4	Family doctor has an after-hour arrangement when office is closed (% yes)	11.6 (8.1 - 15.1)	15.9 (11.2 - 20.6)	15	18.2 (17.5 - 18.9)	
	A-5	How quickly appointment can be made with family doctor	(% on same day or next day)	14.8 (10.9 - 18.7)	22.2 (16.8 - 27.6)	27	30.1 (29.3 - 30.9)
			(% within 5 days)	44.7 (39.1 - 50.4)	47.2 (40.7 - 53.7)	28	60.3 (59.4 - 61.2)
	A-6	Nurse working with family doctor who is regularly involved in health care (% yes)	33.2 (28.0 - 38.3)	31.5 (25.6 - 37.5)	12	27.5 (26.7 - 28.3)	
	A-7	Has access to a primary health team ³ (% yes)	37.6 (32.4 - 42.8)	33.5 (27.6 - 39.4)	11	28.5 (27.8 - 29.2)	
	A-8	Wait time at the hospital emergency department (% less than 4 hours)	71.2 (64.4 - 78.0)	71.0 (63.1 - 78.9)	19	73.9 (72.7 - 75.1)	
A-9	How often citizens received the health care service they needed in the official language of their choice (% who always received service in language of preference)	English ⁴	78.1 (55.6 - 100.0)	75.6^E (47.5 - 100.0)	-	95.2 (94.8 - 95.6)	
		French ⁴	81.5 (77.1 - 85.9)	75.7 (70.1 - 81.3)	-	72.6 (71.0 - 74.2)	

² Indicator rank across 33 communities, from best (1) to worst (33)

³ Primary health team: Has access to a family doctor and has access to either a nurse practitioner or a nurse or other health professionals at their family doctor's office

⁴ Preferred language of service as indicated by respondent in the survey

^E Use with caution (coefficient of variation between 16.7% and 33.3%)

^F Too unreliable to be published (coefficient of variation greater than 33.3%)

Confidence interval calculated at a 95% level of confidence.

Symbols **B** and **W** indicate a statistically significant difference at the 95% level of confidence.

Sample size shown represents total number of respondents. Exact number of respondents varies per survey indicator.

Tracadie-Sheila, Saumarez and Saint-Isidore

B = Better than 2011 results
W = Worse than 2011 results

n=337 n=240 1 to 33 n=13,614

Primary Health Indicator		2011	2014	Rank ²	NB
Appropriateness					
Care/service provided is relevant to the patients'/clients' needs and based on established standards					
Q-1	Adults 65 and over who received their flu shot in the last year (% yes)	Not available	69.8 (56.3 - 83.3)	18	70.6 (69.2 - 72.0)
Q-2	Adults with one or more of four select chronic health conditions ⁵ who had measurements for blood pressure in the past 12 months (% yes)	86.0 (80.3 - 91.6)	83.9 (77.0 - 90.8)	30	91.3 (90.6 - 92.0)
Q-3	Adults with one or more of four select chronic health conditions ⁵ who had measurements for cholesterol in the past 12 months (% yes)	78.1 (71.3 - 84.8)	84.2 (77.3 - 91.1)	3	76.3 (75.2 - 77.4)
Q-4	Adults with one or more of four select chronic health conditions ⁵ who had measurements for blood sugar in the past 12 months (% yes)	73.6 (66.2 - 80.9)	75.6 (67.5 - 83.7)	17	73.9 (72.8 - 75.0)
Q-5	Adults with one or more of four select chronic health conditions ⁵ who had measurements for body weight in the past 12 months (% yes)	61.9 (53.9 - 69.9)	55.9 (46.4 - 65.4)	22	59.3 (58.0 - 60.6)
Q-6	How often family doctor helps citizens coordinate the care from other health care providers and places (% always)	58.7 (52.2 - 65.2)	58.0 (49.9 - 66.1)	32	70.7 (69.7 - 71.7)
Effectiveness					
Care/service, intervention or action achieves the desired results					
Q-7	Citizens with a chronic health condition ⁷ who are confident in controlling and managing their health condition (% very confident)	38.0 (31.7 - 44.2)	35.7 (28.5 - 42.9)	31	42.2 (41.2 - 43.2)
Safety					
Potential risks of an intervention or the environment are avoided or minimized					
Q-8	Citizens with a chronic health condition ⁷ who know what each of their prescribed medications do (% strongly agree)	25.5 (19.5 - 31.5)	26.0 (18.9 - 33.1)	30	47.7 (46.6 - 48.8)
Q-9	Was harmed due to a medical error or mistake as a result of health care services received in the last 12 months (excluding hospital stay) (% yes)	F	4.0 ^{E,6} (1.5 - 6.5)	-	2.7 ⁶ (2.4 - 3.0)

² Indicator rank across 33 communities, from best (1) to worst (33)

⁵ Self-reported by respondent: Diabetes, heart disease, stroke or high blood pressure

⁶ For this type of indicator, a "better" result is achieved when the indicator has a "lower" value

⁷ Self-reported by respondent: Arthritis, asthma, cancer, chronic pain, depression, diabetes, emphysema or COPD, gastric reflux (GERD), heart disease, high blood pressure or hypertension, mood disorder other than depression, or stroke

^E Use with caution (coefficient of variation between 16.7% and 33.3%)

^F Too unreliable to be published (coefficient of variation greater than 33.3%)

Confidence interval calculated at a 95% level of confidence.

Symbols **B** and **W** indicate a statistically significant difference at the 95% level of confidence.

Sample size shown represents total number of respondents. Exact number of respondents varies per survey indicator.

Tracadie-Sheila, Saumarez and Saint-Isidore

B = Better than 2011 results
W = Worse than 2011 results

n=337 n=240 1 to 33 n=13,614

Primary Health Indicator		2011	2014	Rank ²	NB
Efficiency					
Achieving the desired results with the most cost-effective use of resources					
Q-10	Model of care used most often when sick or in need of care from a health professional (% hospital emergency department)	21.4 ⁶ (17.0 - 25.8)	20.8 ⁶ (15.7 - 25.9)	27	11.5 ⁶ (11.0 - 12.0)
Q-11	Has used Tele-Care or other telephone information lines in the last 12 months (% yes)	8.1 ^E (5.2 - 11.0)	5.3 ^E (2.5 - 8.1)	26	7.4 (7.0 - 7.8)
Use of services					
Using primary health services in the last 12 months					
U-1	Visited a personal family doctor (% yes)	80.5 (76.3 - 84.7)	92.4 (89.1 - 95.7)	1	80.3 (79.6 - 81.0)
U-2	Visited a hospital emergency department (% yes)	52.5 (47.2 - 57.8)	61.0 (54.8 - 67.2)	2	41.3 (40.5 - 42.1)
U-3	Visited a specialist (% yes)	Not available	59.5 (53.4 - 65.6)	8	56.4 (55.6 - 57.2)
U-4	Visited an after-hours clinic or a walk-in clinic (% yes)	21.6 (17.2 - 26.0)	16.0 (11.4 - 20.6)	23	24.3 (23.6 - 25.0)
U-5	Visited an alternative practitioner (% yes)	Not available	27.5 (21.9 - 33.1)	9	25.2 (24.5 - 25.9)
U-6	Visited a community health centre (% yes)	6.1 ^E (3.5 - 8.7)	7.6 ^E (4.3 - 10.9)	14	6.9 (6.5 - 7.3)
U-7	Used ambulance services (% yes)	4.7 ^E (2.4 - 7.0)	5.8 ^E (2.9 - 8.7)	14	5.7 (5.3 - 6.1)
U-8	Visited a nurse practitioner (% yes)	7.8 ^E (4.9 - 10.7)	12.3 ^E (8.2 - 16.4)	6	7.7 (7.3 - 8.1)
U-9	Visited a health professional at a private clinic (% yes)	Not available	26.1 (20.6 - 31.6)	8	23.7 (23.0 - 24.4)

² Indicator rank across 33 communities. Under “efficiency” the rank is from best (1) to worst (33). Under “use of services” the rank is from highest (1) to lowest (33).

⁶ For this type of indicator, a “better” result is achieved when the indicator has a “lower” value

^E Use with caution (coefficient of variation between 16.7% and 33.3%)

Confidence interval calculated at a 95% level of confidence.

Symbols **B** and **W** indicate a statistically significant difference at the 95% level of confidence.

Sample size shown represents total number of respondents. Exact number of respondents varies per survey indicator.

Tracadie-Sheila, Saumarez and Saint-Isidore

B = Better than 2011 results
W = Worse than 2011 results

n=337 n=240 1 to 33 n=13,614

Primary Health Indicator		2011	2014	Rank ²	NB
Communication/Relationship with personal family doctor					
C-1	How often family doctor explains things in a way that is easy to understand (% always)	69.1 (63.6 - 74.7)	76.8 (71.3 - 82.3)	24	80.2 (79.5 - 80.9)
C-2	How often family doctor involves citizens in decisions about their health care (% always)	44.9 (39.0 - 50.8)	49.0 (42.4 - 55.6)	32	68.2 (67.3 - 69.1)
C-3	How often family doctor gives citizens enough time to discuss feelings, fears and concerns about their health (% always)	62.1 (56.4 - 67.8)	69.6 (63.0 - 76.2)	26	71.9 (71.0 - 72.8)
Satisfaction (Overall rating of primary health services. When rating is described as “% 8, 9 or 10”, this is based on a scale of zero to 10.)					
S-1	Overall health care services received in New Brunswick (% 8, 9 or 10)	62.2 (57.0 - 67.4)	65.4 (59.4 - 71.4)	23	67.9 (67.1 - 68.7)
S-2	Services from personal family doctor (% 8, 9 or 10)	73.2 (68.1 - 78.3)	81.4 (76.3 - 86.5)	28	83.9 (83.2 - 84.6)
S-3	Services at the hospital emergency department (% 8, 9 or 10)	44.3 (36.9 - 51.7)	52.3 (43.8 - 60.8)	19	53.4 (52.1 - 54.7)
S-4	Services from specialist (% 8, 9 or 10)	Not available	77.3 (70.7 - 83.9)	30	82.0 (81.2 - 82.8)
S-5	Services at an after-hours clinic or walk-in clinic (% 8, 9 or 10)	80.9 (71.8 - 90.0)	84.2 (73.2 - 95.2)	3	73.3 (71.6 - 75.0)
S-6	Services from alternative practitioner (% 8, 9 or 10)	Not available	82.8 (74.1 - 91.5)	29	87.3 (86.2 - 88.4)
S-7	Services at a community health centre (% 8, 9 or 10)	80.2 (63.9 - 96.5)	84.0 (66.6 - 100.0)	10	76.2 (73.6 - 78.8)
S-8	Ambulance services (% 8, 9 or 10)	76.0 (55.1 - 96.9)	74.0 ^E (50.2 - 97.8)	29	87.5 (85.3 - 89.7)
S-9	Services from nurse practitioner (% 8, 9 or 10)	Suppressed ⁸	Suppressed ⁸	-	85.8 (83.7 - 87.9)
S-10	Services at a private clinic (% 8, 9 or 10)	Not available	84.9 (75.1 - 93.9)	14	82.9 (81.6 - 84.2)
S-11	Evaluation of Tele-Care services (% very or somewhat helpful)	86.9 (74.2 - 99.6)	B 100.0 (100 - 100)	1	89.3 (87.2 - 91.4)
S-12	Evaluation of care received for mental or emotional health (among citizens who have talked to a health professional) (% very or somewhat helpful)	84.9 (75.4 - 94.4)	B 98.4 (94.6 - 100.0)	2	90.8 (89.6 - 92.0)
S-13	Wait time from booking appointment to seeing specialist (among citizens who have seen a specialist) (% very or somewhat satisfied)	Not available	83.8 (78.0 - 89.6)	8	80.5 (79.6 - 81.4)

² Indicator rank across 33 communities, from best (1) to worst (33)

^F Too unreliable to be published (coefficient of variation greater than 33.3%)

⁸ Results suppressed due to limited number of nurse practitioners

^E Use with caution (coefficient of variation between 16.7% and 33.3%)

Sample size shown represents total number of respondents.
 Exact number of respondents varies per survey indicator.

Confidence interval calculated at a 95% level of confidence.

Symbols **B** and **W** indicate a statistically significant difference at the 95% level of confidence.